


ISSN 2072-5981



***agnetic
Resonance
in Solids***

Electronic Journal

*Volume 19,
Issue 2
Paper No 17201,
1-11 pages
2017*

<http://mrsej.kpfu.ru>

<http://mrsej.ksu.ru>



Established and published by Kazan University
Sponsored by International Society of Magnetic Resonance (ISMAR)
Registered by Russian Federation Committee on Press, August 2, 1996
First Issue was appeared at July 25, 1997

© Kazan Federal University (KFU)*

"Magnetic Resonance in Solids. Electronic Journal" (MRSej) is a peer-reviewed, all electronic journal, publishing articles which meet the highest standards of scientific quality in the field of basic research of a magnetic resonance in solids and related phenomena.

Indexed and abstracted by
Web of Science (ESCI, Clarivate Analytics, from 2015), Scopus (Elsevier, from 2012), RusIndexSC (eLibrary, from 2006), Google Scholar, DOAJ, ROAD, CyberLeninka (from 2006), SCImago Journal & Country Rank, etc.

Editors-in-Chief

Jean Jeener (Universite Libre de Bruxelles, Brussels)

Boris Kochelaev (KFU, Kazan)

Raymond Orbach (University of California, Riverside)

Executive Editor

Yurii Proshin (KFU, Kazan)

mrsej@kpfu.ru

Editors

Vadim Atsarkin (Institute of Radio Engineering and Electronics, Moscow)

Yurij Bunkov (CNRS, Grenoble)

Mikhail Eremin (KFU, Kazan)

David Fushman (University of Maryland, College Park)

Hugo Keller (University of Zürich, Zürich)

Yoshio Kitaoka (Osaka University, Osaka)

Boris Malkin (KFU, Kazan)

Alexander Shengelaya (Tbilisi State University, Tbilisi)

Jörg Sichelschmidt (Max Planck Institute for Chemical Physics of Solids, Dresden)

Haruhiko Suzuki (Kanazawa University, Kanazawa)

Murat Tagirov (KFU, Kazan)

Dmitrii Tayurskii (KFU, Kazan)

Valentine Zhikharev (KNRTU, Kazan)



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).



This is an open access journal which means that all content is freely available without charge to the user or his/her institution. This is in accordance with the [BOAI definition of open access](https://www.boai.gov.ru/).

* In Kazan University the Electron Paramagnetic Resonance (EPR) was discovered by Zavoisky E.K. in 1944.

Random walks in disordered lattice, CTRW, memory and dipole transport

F.S. Dzheparov

National Research Center "Kurchatov Institute", ITEP, Moscow 117258, Russia

National Research Nuclear University "MEPhI", Moscow 115409, Russia

E-mail: dzheparov@itep.ru

(Received October 30, 2017; revised November 11, 2017; accepted November 14, 2017)

Application of CTRW (continuous time random walks) to dipole hopping transport is reviewed. Conditions of applicability of basic kinetic equations to spin systems are indicated. Correct versions of derivation of the CTRW-equations are presented. Existence of different forms of memory kernels is demonstrated. Correction of Scher-Lax memory kernel within geometrical memory approach is fulfilled in accordance with leading terms of concentration expansion. Approximate solution for autocorrelation function is considered. Modern state of numerical simulation and experimental measurements of autocorrelation function in nuclear polarization delocalization are described. It is shown, that application of the CTRW was more successful in description of dipole transport than for hopping conductivity.

PACS: 45.10.Hj, 76.60.-k, 82.56.-b, 76.30.-v, 05.40.-a

Keywords: random walks, random media, disordered lattice, survival probability, dipole transport, hopping transport, master equation, memory kernel, projection operator, spin diffusion

1. Introduction

Random walks in disordered lattice are described by the equations

$$\frac{\partial}{\partial t} p_{i0} = -\sum_j (w_{ji} p_{i0} - w_{ij} p_{j0}), \quad p_{i0}(t=0) = \delta_{i0}, \quad (1)$$

where $p_{i0}(t)$ is the "probability" to find an "excitation" at position \mathbf{r}_i , if it started at $\mathbf{r}_0 = \mathbf{0}$ at $t=0$ and δ_{ij} is the Kronecker's symbol. Here w_{ji} is the transition rate for transfer from \mathbf{r}_i to \mathbf{r}_j . The problem of disordered sites will be considered below, when positions \mathbf{r}_j are statically and randomly distributed on sites of regular lattice, w_{ji} depends on $\mathbf{r}_j = \mathbf{r}_i - \mathbf{r}_j$, and observables are directly related to solution $p_{i0}(t)$, averaged over all possible positions $\{\mathbf{r}_j\}$. Occupation number representation allows to rewrite Eqs.(1) as [1]

$$\frac{\partial}{\partial t} \tilde{P}_{\mathbf{x}0} = -\sum_z n_z n_z (w_{z\mathbf{x}} \tilde{P}_{\mathbf{x}0} - w_{\mathbf{x}z} \tilde{P}_{z0}) = -\left(A^{(0)} \tilde{P}\right)_{\mathbf{x}0}, \quad (2)$$

or

$$\frac{\partial}{\partial t} \tilde{P}_{\mathbf{x}0} = -\sum_z (n_z w_{z\mathbf{x}} \tilde{P}_{\mathbf{x}0} - n_x w_{\mathbf{x}z} \tilde{P}_{z0}) = -\left(A^{(1)} \tilde{P}\right)_{\mathbf{x}0}, \quad (3)$$

where the propagator $\tilde{P}_{\mathbf{x}0}(t)$ gives the probability to find the excitation at lattice site \mathbf{x} , when initially it was at the site $\mathbf{0}$, and $w_{z\mathbf{x}} = w_{ij}(r_i = \mathbf{x}, r_j = \mathbf{z})$. Here n_r is occupation number of the site \mathbf{r} by a donor ($n_r = 1(0)$ if the site \mathbf{r} is (not) occupied by the donor), while the donor is an impurity, which can carry the excitation. Eqs. (3) are equivalent to Eqs. (2) because $n_x \tilde{P}_{\mathbf{x}0} = \tilde{P}_{\mathbf{x}0}$, while equivalence of Eqs. (1) and (2) is evident, if we omit in (2) all empty sites for which $n_x = 0$ and, consequently, $\tilde{P}_{\mathbf{x}0} = 0$.

The problem consists in calculation of the observable propagator

$$P_{\mathbf{x}0}(t) = \left\langle \tilde{P}_{\mathbf{x}0}(t) \right\rangle. \quad (4)$$

Here $\langle \dots \rangle$ is averaging over all possible donors positions in infinite sample.

Solutions to Eqs. (2) and (3) are of the form $\tilde{P}_{\mathbf{x}\mathbf{0}} = (\exp(-At))_{\mathbf{x}\mathbf{0}} n_{\mathbf{0}} / c$, where the operator $A = A^{(0)}$ or $A = A^{(1)}$ and $c = \langle n_{\mathbf{x}} \rangle$ is the probability of occupation (concentration). The initial condition

$$\tilde{P}_{\mathbf{x}\mathbf{0}}(t=0) = n_{\mathbf{x}} \delta_{\mathbf{x}\mathbf{0}} / c \quad (5)$$

ensures, that the excitation can be placed at site $\mathbf{0}$ if it is occupied by a donor, and produces evident normalization

$$\sum_{\mathbf{x}} P_{\mathbf{x}\mathbf{0}}(t) = 1.$$

Occupations of different sites are assumed as independent and having no dependence on the \mathbf{x} (with small $c \ll 1$ as a rule):

$$\langle n_{\mathbf{r}} \rangle = c, \quad \langle n_{\mathbf{r}} n_{\mathbf{x}} \rangle = c \delta_{\mathbf{r}\mathbf{x}} + c^2 (1 - \delta_{\mathbf{r}\mathbf{x}}), \quad \langle \prod_{j=1}^m n_{\mathbf{r}_j} \rangle = \prod_{j=1}^m \langle n_{\mathbf{r}_j} \rangle = c^m.$$

All \mathbf{r}_j are different in the last relation. Coincidence in indexes can be treated using the identity $n_{\mathbf{r}}^2 = n_{\mathbf{r}}$.

Most important experimental realizations of the process (1) are hopping conductivity, for which the CTRW theory of Scher and Lax [2] was developed, Förster electronic energy transfer [3] and spin-polarization transfer [1]. The simplest transition rates are of the form

$$w_{\mathbf{z}\neq\mathbf{x}} = w_0 \exp(-|\mathbf{z} - \mathbf{x}| / r_0), \quad w_{\mathbf{x}\mathbf{x}} = 0 \quad (6)$$

for conductivity problem, and

$$w_{\mathbf{z}\neq\mathbf{x}} = w_0 r_0^6 / (\mathbf{z} - \mathbf{x})^6, \quad w_{\mathbf{x}\mathbf{x}} = 0 \quad (7)$$

for dipole transitions in electronic energy and spin-polarization transfers. Here w_0 corresponds to the transition rate at distance r_0 between donors. More advanced representations for transition rates can be found in special literature, see, for example, [4] for spin polarization transfer. It should be noted that in the problem of spin-polarization transfers the propagator $P_{\mathbf{x}\mathbf{0}}(t)$ represents polarization (instead of probability) of a spin at the site \mathbf{x} , when initially polarization was localized at the site $\mathbf{0}$.

Polarization transport in disordered media is very important in spin kinetics and magnetic resonance. Its manifestations are related, first of all, with spin-lattice and spin-spin relaxation [5-7] as well as with establishing and retaining of (quasi)equilibrium [8]. The discovery of the model system $^8\text{Li}-^6\text{Li}$ is of great importance, because it gives a unique possibility for direct measurement of the survival probability $P_{\mathbf{0}\mathbf{0}}(t)$ (see [4] and reference therein) contrary to other systems, where the process (1) can not be directly observable.

The conditions of applicability of Eqs. (1-3) to description of spin polarization transport in the subsystem of impurity spins require that the speed of the process should be small a) relative to the speed of phase relaxation of impurity spins and b) in comparison with the speed of fluctuations of local fields on these spins [1, 4]. The model system $^8\text{Li}-^6\text{Li}$ in the LiF single crystal represents excellent example for experimental study of the problem. It consists of one beta-active nucleus ^8Li with high initial polarization and of ^6Li nuclei with negligible thermal initial polarization. The nuclei ^6Li have small controllable concentration, and the difference of gyromagnetic ratios of ^6Li and ^8Li is rather small: $(g_8 - g_6) / g_6 = 0.0057$. The matrix LiF consists of nuclei ^7Li and ^{19}F with higher gyromagnetic ratios that ensures with high quality the conditions of applicability, mentioned above, in comfortable external magnetic field in the range $200\text{G} < H_0 < 3\text{kG}$ [4]. The evolution of the polarization $P_{\mathbf{0}\mathbf{0}}(t)$ of the nuclei ^8Li is directly observable via measurement of the beta-decay asymmetry (β -NMR).

We use here the simplest transition rates (7) for the problem of spin polarization transport in order to concentrate our attention on specifics of random walks in disordered media. Quantitative comparison with experimental results requires to take into account that transition rate $w_{\mathbf{z}\mathbf{x}}$ depends as well on the orientation of $\mathbf{x} - \mathbf{z}$ relative to external static magnetic field, on correlation of local fields on donor spins and on difference in spin values of ^8Li and ^6Li [4].

The problems of calculation of the propagator (4) or memory kernels $N_{\mathbf{z}\mathbf{x}}(t)$ in corresponding master equation

$$\frac{\partial}{\partial t} P_{\mathbf{x}\mathbf{0}}(t) = - \int_0^t dt' \sum_{\mathbf{z}} [N_{\mathbf{z}\mathbf{x}}(t') P_{\mathbf{x}\mathbf{0}}(t-t') - N_{\mathbf{x}\mathbf{z}}(t') P_{\mathbf{z}\mathbf{0}}(t-t')], \quad P_{\mathbf{x}\mathbf{0}}(t=0) = \delta_{\mathbf{x}\mathbf{0}} \quad (8)$$

belong to the most complex problems of modern statistical mechanics and they have not adequate analytical solution up to now. Nonseparable version of CTRW theory, developed by Scher and Lax in two articles [2], produced an integral representation for approximate solution of the problem and extended previously developed separable CTRW of Montroll and Weiss [9] (which operated with kernels of the form: $N_{\mathbf{z}\mathbf{x}}(t) = X_{\mathbf{z}\mathbf{x}} \cdot Y_{\mathbf{x}}(t)$).

CTRW is the abbreviation of continuous time random walks. The meaning of the expression consists in following. If the excitation is placed at site \mathbf{x} , then its escape, according to Eqs. (1)-(3), is described by the law $\tilde{Q}_{\mathbf{x}}(t) = \exp(-t/\tau_{\mathbf{x}})$, with one characteristic hopping time $\tau_{\mathbf{x}} = \left(\sum_{\mathbf{z}} n_{\mathbf{z}} w_{\mathbf{z}\mathbf{x}}\right)^{-1}$, while the Eq. (8) corresponds to the escape law with $dQ_{\mathbf{x}}(t)/dt = -\int_0^t dt' \sum_{\mathbf{z}} N_{\mathbf{z}\mathbf{x}}(t') Q_{\mathbf{x}}(t-t')$ and its result can be written as $Q_{\mathbf{x}}(t) = \int d\tau \exp(-t/\tau) W_{\mathbf{x}}(\tau)$ with continuous distribution of times $W_{\mathbf{x}}(\tau)$.

The Scher-Lax theory was directed on calculation of frequency dependent diffusion (or conductivity), which are defined by

$$\langle \mathbf{x}^2(t) \rangle = \sum_{\mathbf{x}} \mathbf{x}^2 P_{\mathbf{x}\mathbf{0}}(t),$$

and it produced important progress in the field at that time. But the theory was ineffective in description of other important quantity, survival probability (or autocorrelation function) $P_{\mathbf{0}\mathbf{0}}(t)$, which is directly measurable in fine optical [10, 11] and beta-NMR [4, 12] studies. Other important property - conceptual foundation of the relations between “microscopic” equations (1)-(3) and master equations (8) is still unrecognized by absolute majority of workers, that is clearly seen from legend about the derivation of the master equation which exists 37 years and passed in erroneous form through many reviews, see for example [13, ch.5] and [14].

The aims of this article consists in explaining of nonseparable version of CTRW, constructed by me and my colleagues for qualitatively correct description of the autocorrelator $P_{\mathbf{0}\mathbf{0}}(t)$ in problems of dipole hopping transport, in improvement of some details of the theory, in short description of corresponding modern numerical and experimental results and in concentrated clarifying of those connections of the master equations (8) with primary equations (1)-(3), which are fundamentally important for correct calculation of the $P_{\mathbf{0}\mathbf{0}}(t)$ and for the problem of random walks on disordered sites as a whole. We resume our experience of construction and application of the CTRW theory in the Section 6 with some differences relative to “incredible possibilities of the CTRWs” indicated in Ref. [14].

2. Reformulation and generalization of the CTRW theory

The method of approximate construction of the propagator $P_{\mathbf{x}\mathbf{0}}(t)$, developed in Ref. [2], was reformulated in [15, 16] (see also [17, 18] for different approach) basing on the following assumptions:

a) the first term in square brackets in (8) is the rate of polarization outflow from \mathbf{x} to \mathbf{z} , while the second term is the rate of inflow from \mathbf{z} to \mathbf{x} ;

b) inflow rate from \mathbf{z} to \mathbf{x} does not depend on how the polarization reached \mathbf{z} .

The processes of inflow to the site and outflow from it are now separated, and to determine the kernels $N_{\mathbf{zx}}(t)$ we can treat the simple case in which an exact averaging can be carried out. For this purpose, we choose the process of the outflow of polarization from an arbitrary site in exactly solvable problem

$$\begin{aligned} \frac{\partial}{\partial t} \tilde{F}_{\mathbf{xx}} &= -\sum_{\mathbf{z}} n_{\mathbf{z}} w_{\mathbf{zx}} \tilde{F}_{\mathbf{zx}}, & \frac{\partial}{\partial t} \tilde{F}_{\mathbf{z\neq x}} &= n_{\mathbf{z}} w_{\mathbf{zx}} \tilde{F}_{\mathbf{zx}}, \\ \tilde{F}_{\mathbf{zx}}(t=t_0) &= n_{\mathbf{z}} \delta_{\mathbf{zx}} / c, & \tilde{F}_{\mathbf{zx}}(t < t_0) &= 0. \end{aligned} \quad (9)$$

According to the assumptions above, the average $F_{\mathbf{zx}}(t) = \langle \tilde{F}_{\mathbf{zx}}(t) \rangle$ must satisfy the equations

$$\begin{aligned} \frac{\partial}{\partial t} F_{\mathbf{xx}}(t) &= -\int_0^t dt' \sum_{\mathbf{z}} N_{\mathbf{zx}}(t') F_{\mathbf{zx}}(t-t'), & \frac{\partial}{\partial t} F_{\mathbf{z\neq x}}(t) &= \int_0^t dt' N_{\mathbf{zx}}(t') F_{\mathbf{zx}}(t-t'), \\ F_{\mathbf{zx}}(t=t_0) &= \delta_{\mathbf{zx}}, & F_{\mathbf{zx}}(t < t_0) &= 0. \end{aligned} \quad (10)$$

As a result, the equation for the kernel $N_{\mathbf{zx}}(t)$ obtains the form

$$\int_0^t dt' N_{\mathbf{zx}}(t') \left\langle \exp\left(-\sum_{\mathbf{q}} n_{\mathbf{q}} w_{\mathbf{qx}}(t-t')\right) \right\rangle = \left\langle n_{\mathbf{z}} w_{\mathbf{zx}} \exp\left(-\sum_{\mathbf{q}} n_{\mathbf{q}} w_{\mathbf{qx}} t\right) \right\rangle \quad (11)$$

with

$$\left\langle \exp\left(-\sum_{\mathbf{q}} n_{\mathbf{q}} w_{\mathbf{qx}} t\right) \right\rangle = \exp\left(\sum_{\mathbf{q}} \ln\left(1 - c\left(1 - e^{-w_{\mathbf{qx}} t}\right)\right)\right), \quad (12)$$

$$\left\langle n_{\mathbf{z}} w_{\mathbf{zx}} \exp\left(-\sum_{\mathbf{q}} n_{\mathbf{q}} w_{\mathbf{qx}} t\right) \right\rangle = c w_{\mathbf{zx}} e^{-w_{\mathbf{zx}} t} \exp\left(\sum_{\mathbf{q} \neq \mathbf{z}} \ln\left(1 - c\left(1 - e^{-w_{\mathbf{qx}} t}\right)\right)\right). \quad (13)$$

The Eqs. (8) and (11) give qualitatively satisfactory description of those properties of the hopping conductivity and delocalization of excitations, which are related with $\langle \mathbf{x}^2(t) \rangle = \sum_{\mathbf{x}} \mathbf{x}^2 P_{\mathbf{x0}}(t)$, but, as it is demonstrated in the next section, they are erroneous in the description of the autocorrelator $P_{\mathbf{00}}(t)$, which is directly measurable in the optical [3, 10, 11] and beta-NMR [4, 12] experiments.

3. Main weakness of the Scher-Lax CTRW theory

In order to sharpen the problem we can consider the continuum media approximation, when impurity concentration $c \rightarrow 0$ and lattice prime cell volume $\Omega \rightarrow 0$ at a fixed value of impurity density $n = c / \Omega$. In the continuum media approximation the Eqs. (8) obtain the form

$$\begin{aligned} \frac{\partial}{\partial t} P(\mathbf{x}, t | \mathbf{0}) &= -\int_0^t dt' \int d^3 z \left[N_{\mathbf{zx}}(t') P(\mathbf{x}, t-t' | \mathbf{0}) - N_{\mathbf{zx}}(t') P(\mathbf{z}, t-t' | \mathbf{0}) \right] = \\ &= -\int_0^t dt' \int d^3 z Z_{\mathbf{zx}}(t') P(\mathbf{z}, t-t' | \mathbf{0}), \quad P(\mathbf{z}, t=0 | \mathbf{0}) = \delta(\mathbf{x}), \end{aligned} \quad (14)$$

where $\delta(\mathbf{x})$ is Dirac's delta-function, the probability density $P(\mathbf{x}, t | \mathbf{0}) = P_{\mathbf{x0}}(t) / \Omega$ has normalization $\int d^3 x P(\mathbf{x}, t | \mathbf{0}) = 1$ and the kernels can be written as

$$N_{\mathbf{zx}}(\lambda) = \int_0^\infty dt e^{-\lambda t} N_{\mathbf{zx}}(t) = n w_{\mathbf{zx}} Q(\lambda + w_{\mathbf{zx}}) / Q(\lambda), \quad (15)$$

$$Q(\lambda) = \int_0^\infty dt e^{-\lambda t} Q(t) = \left(\lambda + \int d^3 z N_{\mathbf{zx}}(\lambda) \right)^{-1},$$

$$Q(t) = \left\langle \exp\left(-\sum_{\mathbf{q}} n_{\mathbf{q}} w_{\mathbf{q}\mathbf{x}} t\right) \right\rangle = \exp\left(-n \int d^3 q \left(1 - e^{-w_{\mathbf{q}\mathbf{x}} t}\right)\right) = \exp\left(-(\beta t)^{1/2}\right). \quad (16)$$

Here and below we apply the same symbols for time dependent functions and for their Laplace transformations distinguishing them by the argument. The Förster constant $\beta = (16/9)\pi^3 n^2 w_0 r_0^6$ is defined by the last equality in (16).

The master equation (14) can be rewritten in the Laplace representation as

$$P(\mathbf{x}, \lambda | \mathbf{0}) = Q(\lambda) \left(\delta(\mathbf{x}) + \int d^3 z N_{\mathbf{z}\mathbf{x}}(\lambda) P(\mathbf{z}, \lambda | \mathbf{0}) \right) = P^{(s)}(\mathbf{x}, \lambda | \mathbf{0}) + P^{(r)}(\mathbf{x}, \lambda | \mathbf{0}), \quad (17)$$

or, in the time representation

$$P(\mathbf{x}, t | \mathbf{0}) = P^{(s)}(\mathbf{x}, t | \mathbf{0}) + P^{(r)}(\mathbf{x}, t | \mathbf{0}), \quad P^{(s)}(\mathbf{x}, t | \mathbf{0}) = Q(t) \delta(\mathbf{x}). \quad (18)$$

The solution (15) indicates that $N_{\mathbf{z}\mathbf{x}}(\lambda)$ is a smooth function of $|\mathbf{x} - \mathbf{z}|$, therefore Eqs. (17) and (18) separate the propagator on singular $P^{(s)}(\mathbf{x}, t | \mathbf{0})$ and regular $P^{(r)}(\mathbf{x}, t | \mathbf{0})$ parts near $x = 0$. At that, the singular part is defined only by the singular part of the memory kernel $Z_{\mathbf{z}\mathbf{x}}^{(s)}(t) = \delta(\mathbf{x} - \mathbf{z}) \int d^3 q N_{\mathbf{q}\mathbf{x}}(t)$ of the Eq. (14).

Simple analysis indicates, that for long time

$$\langle \mathbf{x}^2(t) \rangle = \int d^3 x x^2 P(\mathbf{x}, t | \mathbf{0}) \sim n^{-2/3} \beta t, \quad (19)$$

that is in agreement with scaling arguments [1, 19] and expected for diffusion long time asymptotics. But the solution $P^{(s)}(\mathbf{x}, t | \mathbf{0}) = F_0(t) \delta(\mathbf{x})$ with $F_0(t) = Q(t)$ is incorrect [1, 11, 15, 16]. It decays with time exponentially, while more slow behavior, diffusion like as $F_0(t) \sim (\beta t)^{-3/2}$ should be expected, because $F_0(t)$ is the survival probability, and it should be of the order of probability $F_1(t)$ to find the excitation on a donor, placed near the origin, that is

$$F_1(t \rightarrow \infty) \sim \int d^3 x \mathcal{G}(x < n^{-1/3}) P_{\mathbf{x}\mathbf{0}}^{(r)}(t \rightarrow \infty) \sim \frac{1}{n} \cdot \frac{1}{\langle x^2(t) \rangle^{3/2}} \sim (\beta t)^{-3/2}. \quad (20)$$

Here Heaviside's function $\mathcal{G}(x)$ is applied.

The relations (17) and (18) indicate, that correct $F_0(t \rightarrow \infty)$ can be obtained if the singular part of the memory kernel $Z_{\mathbf{z}\mathbf{x}}^{(s)}(t) = \delta(\mathbf{x} - \mathbf{z}) Z_D(t)$ has correct long time tail $Z_D(t \rightarrow \infty) = -F_0(t) / \int_0^\infty dt' F_0(t') \sim -(\beta t)^{-3/2}$ [16]. This conclusion contradicts to main strategy of application of the memory functions method when reasonable approximation for short-term memory kernels produces satisfactory long time behavior for the solution of master equation (that was fulfilled in the relation (19) for example). Therefore, we should look for justification of the applicability of the memory function method (i.e. Eq. (8)) and for modification of the memory kernels.

4. Correction of the CTRW theory

The justification of the applicability of the memory function method can be based on derivation of the Eq. (8) applying the Nakajima-Zwanzig projection operator technique to Eqs. (2) or (3). Similar attempt was undertaken in Ref. [20] for Eqs. (2) choosing the projection operator $\hat{\pi}$ as simple averaging $\hat{\pi} A = \langle A \rangle$ for any A . But initial condition was applied in incorrect form $\tilde{P}_{\mathbf{x}\mathbf{0}}(t=0) = \delta_{\mathbf{x}\mathbf{0}}$, therefore all equations, derived in [20], are incorrect [21]. Nevertheless, authors of reviews [13] and [14], as well as more than three hundreds other workers insist that the Ref. [20] gave convincing derivation of the master equation (8) ignoring the criticism of the Ref. [21]. It should be noted

nevertheless that similar derivation can be fulfilled both for Eqs. (2) and (3) with correct initial condition $\tilde{P}_{\mathbf{x}0}(t=0) = n_{\mathbf{x}}\delta_{\mathbf{x}0} / c$ applying other projection operator

$$\hat{\pi} A_{\mathbf{x}} = \frac{n_{\mathbf{x}}}{c} \langle A_{\mathbf{x}} \rangle. \quad (21)$$

Unfortunately, the results of these derivations produce no indication on the way to improve the singular part of memory kernels.

More constructive approach was realized in Refs. [15, 16, 21], see also [22]. Correct initial condition $\tilde{P}_{\mathbf{x}0}(t=0) = n_{\mathbf{x}}\delta_{\mathbf{x}0} / c$ separates only those solutions of the Eqs. (2) and (3), for which the lattice site $\mathbf{r} = \mathbf{0}$ is occupied by a donor, because the identity $n_{\mathbf{x}}R(n_{\mathbf{x}}) = n_{\mathbf{x}}R(1)$ is valid for any reasonable function $R(n_{\mathbf{x}})$. Therefore,

$$P_{\mathbf{x}0}(t) = \left\langle \left(\exp(-At) \right)_{\mathbf{x}0} \frac{n_0}{c} \right\rangle = \left\langle \left(\exp(-At) \right)_{\mathbf{x}0} \right\rangle_0, \quad (22)$$

where $\langle \dots \rangle_0$ means averaging over occupation numbers with the condition $n_0 = 1$. Therefore, we can apply the projector $\hat{\pi} = \pi_0$ acting as

$$\pi_0 B = \langle B \rangle_0 \quad (23)$$

for any B . Standard transformations produce the Eqs. (8) and (14) again, but the memory kernel $N_{\mathbf{z}\mathbf{x}}(t) = N_{\mathbf{z}\mathbf{x}}^{(0)}(t)$ depends both on $\mathbf{z} - \mathbf{x}$ and $\mathbf{x} = \mathbf{x} - \mathbf{0}$, while the propagator $P_{\mathbf{x}0}(t)$ depends on \mathbf{x} only. This new type of memory (geometrical, contrary to dynamical one, which depends on $\mathbf{z} - \mathbf{x}$ only) is much less comfortable for calculations, because the matrix $N_{\mathbf{z}\mathbf{x}}^{(0)}(t)$ can not be diagonalized by the Fourier transformation. Nevertheless [15, 16], equations for the memory kernels can be constructed following the derivation (9)-(13) with the result:

$$\int_0^t dt' N_{\mathbf{z}\mathbf{x}}^{(0)}(t') Q_{\mathbf{x}}^0(t-t') = \frac{c w_{\mathbf{z}\mathbf{x}} \exp(-w_{\mathbf{z}\mathbf{x}} t)}{1 + c(\exp(-w_{\mathbf{z}\mathbf{x}} t) - 1)} Q_{\mathbf{x}}^0(t), \quad \mathbf{z} \neq \mathbf{0}, \quad (24)$$

$$N_{\mathbf{0}\mathbf{x}}^{(0)}(t) = \frac{w_{\mathbf{0}\mathbf{x}}}{c w_{\mathbf{x}\mathbf{0}}} N_{\mathbf{x}\mathbf{0}}^{(0)}(t), \quad Q_{\mathbf{x}}^0(t) = \left\langle \exp\left(-\sum_{\mathbf{z} \neq \mathbf{0}} w_{\mathbf{z}\mathbf{x}} t\right) \right\rangle = \prod_{\mathbf{z} \neq \mathbf{0}} \left(1 + c(e^{-w_{\mathbf{z}\mathbf{x}} t} - 1)\right).$$

Details of the derivation (with taking into account additional exactly solvable model) can be found in Ref. [16].

As a result, the Eqs. (8) with short-term memory (24) produce qualitatively well-formed solution $P_{\mathbf{x}0}(t)$ for all \mathbf{x} and t , and the solution is correct up to terms $\sim c^1$ [15, 16]. It should be noted, that last property was not fulfilled in [2].

5. Approximate solution to corrected CTRW equations

Analytical solution of the Eqs. (8) with kernels (24) is absent. Therefore approximate solution for $P_{\mathbf{0}\mathbf{0}}(t)$ was constructed by matching the short- and long-time asymptotics. Analysis of long-time asymptotics [15, 16] indicates, that for dipole transport

$$P_{\mathbf{0}\mathbf{0}}(\beta t \rightarrow \infty) = \frac{1}{c} G_{\mathbf{0}\mathbf{0}}(t) \left(1 + O\left((\beta t)^{-1}\right)\right), \quad (25)$$

where $G_{\mathbf{x}0}(t)$ is the solution of Eqs. (8) with kernels (11). It can be obtained using the lattice Fourier transformation. For $\beta t \rightarrow \infty$

$$G_{00}(t) = \frac{\Omega}{(2\pi)^3} \int_B d^3k \exp(-N(\mathbf{0})t) \left(1 + O((\beta t)^{-1})\right), \quad (26)$$

where integration is limited by the Brillouin zone and $N(\mathbf{k}) = \sum_{\mathbf{x}} e^{-i\mathbf{k}\mathbf{x}} \int_0^\infty dt N_{\mathbf{x}0}(t)$. The asymptotics $G_{00}(t \rightarrow \infty)$, according to the Laplace's method, is defined by $N(\mathbf{k} \rightarrow 0)$. In simple lattice with cubic symmetry and for dipole transition rates (7) we have

$$N(\mathbf{0}) - N(\mathbf{k}) = Dk^2 - \sigma k^3 + O(k^4). \quad (27)$$

Simple derivation of this expansion can be found in Ref. [23].

The diffusion coefficient D is model dependent, and its value for Scher-Lax theory in continuum media approximation is

$$D_{SL} = n^{4/3} w_0 r_0^6 \frac{\pi^{3/2} 2^{2/3}}{3^{7/3}} \Gamma(1/6) \Gamma(5/3) = \frac{\kappa_{SL}}{6} \beta n^{-2/3}, \quad (28)$$

where $\kappa_{SL} = 0.3725$, while $\sigma = \frac{\pi^2 n}{12} w_0 r_0^6$ is model independent and it is defined by the dipole long ranging exclusively. For example, if we will assume in Eqs. (15) and (16) that $Q_a(t) = \exp(-(\gamma t)^\alpha)$ with arbitrary $\alpha > 0$ and $\gamma > 0$ instead of $Q(t) = \exp(-(\beta t)^{1/2})$, prescribed by the relation (16), then we will obtain other value for the diffusion coefficient but the same $\sigma = \frac{\pi^2 n}{12} w_0 r_0^6$. As a result, in leading terms,

$$P_{00}(\beta t \rightarrow \infty) = \frac{1}{n(4\pi D t)^{3/2}} \left(1 + \frac{4\sigma}{D^{3/2} (\pi t)^{1/2}}\right) = \frac{1}{(\mu\beta t)^{3/2}} \left(1 + \frac{\varphi}{(\mu\beta t)^{1/2}}\right), \quad (29)$$

where $\mu = 0.7801$ and $\varphi = 1.923$.

Short time asymptotics was obtained in Refs. [1, 3, 21, 24]:

$$P_{00}(\beta t < 1) = 1 - (\beta t / 2)^{1/2} + O(\beta t), \quad (30)$$

and the next term $O(\beta t) = d\beta t + O((\beta t)^{3/2})$ was defined in [21] by calculation of the coefficient d basing on exact expansion of the propagator in powers c^m of the concentration.

These results allow to construct the approximation

$$P_{00}(t) = Q(t) + \frac{1 - Q(t)}{(\mu\beta(t + \tau))^{3/2}} \left(1 + \frac{\varphi}{(\mu\beta(t + \tau))^{1/2}}\right), \quad (31)$$

which reproduces the relation (30) up to $(\beta t)^{1/2}$ and both terms of the asymptotics (29) with $\mu\beta\tau = 3.613$ [11, 16].

The coefficients σ and φ prescribe specific evolution of $P_{00}(t)$ relative to its long time asymptote $P_a(t) = (\mu\beta t)^{-3/2}$. At the beginning $P_{00}(t) < P_a(t)$, but with increasing of time we have opposite relation and at $\beta t \rightarrow \infty$ they coincide. This property (reoscillation) was applied in the Ref. [11] to clarify, that, in agreement with Eq. (31), the onset of the diffusive asymptotic behavior in the kinetics of the electrodipole delocalization of excitations in a disordered system of donors takes place at $P_{00}(t) < 0.03$.

Consequent numerical [25, 26] and experimental (beta-NMR) [27, 28] studies revealed, that

a) for small concentrations (i.e. for strongest disorder) the diffusion coefficient is $D_0 = \frac{\kappa_0}{6} \beta n^{-2/3}$ with $\kappa_0=0.296$, this is not very far from $\kappa_{SL} = 0.3725$ in the relation (28), and

b) for $c \leq 0.1$ and for all t autocorrelator $P_{00}(t)$ is of the form

$$P_{00}(t) = F_{00}(t)(1 + \chi(t)), \quad (32)$$

where $F_{00}(t)$ is defined by the relation (31) with more correct diffusion tensor (or coefficient, for isotropic transfer) and variation $\chi(t)$ is relatively small ($-0.1 < \chi(t) < 1$). The variation increases the reoscillation. It should not contain the short- and long-time asymptotic terms, included in the relation (31).

According to the relation (29), for recalculation to new value of the diffusion coefficient, the relation (31) can be written as

$$F_{00}(t) = Q(t) + \frac{1 - Q(t)}{(\mu\beta(t + \tau))^{3/2}} \left(1 + \frac{D_{SL}}{D} \cdot \frac{\varphi}{(\mu\beta(t + \tau))^{1/2}} \right) \quad (33)$$

in order to apply the same value φ . The multiplier D_{SL} / D was forgotten in preceding studies, but it did not produced errors in description of results of numerical studies, because 1) for approximations of the numerical results the relation (32) was applied as a whole with the fitting function $\chi(t)$, and corresponding errors was compensated by $\chi(t)$, and 2) for studied values of D with corresponding $\mu\beta\tau$ the relative inaccuracy in calculation of $F_{00}(t)$ never exceeded 0.05. Nevertheless we should expect that the correction will become important with increasing of accuracy of theoretical and experimental studies, because it allows to exclude the term $\sim (\beta t)^{-1/2}$ from the fitting function $\chi(t)$.

6. Conclusions

As a whole we see, that the version of CTRW, invented in Ref. [2] and improved in Refs. [15, 16], produced important part of the basis for consequent quantitative understanding and experimental investigations of the problem of random walks in disordered media with dipole transitions. It should be noted, that this approach allowed for the first time to obtain analytically correct diffusion long-time asymptotics for autocorrelator $P_{00}(t) \sim (\beta t)^{-3/2}$, while other methods (see, for example, Refs. [3, 24, 29]) produced reasonable behavior at $\beta t \sim 1$, but exponential long time tail. Diffusion long time tail can be obtained for coarse-grained propagator, as in the Ref. [30], this is evident from the relation (20), but it produces no direct information for comparison with precise optical and beta-NMR studies.

We can state, that the CTRW theory was more successful in the description of dipole processes, than for hopping conductivity, where the percolation theory is more applicable. Indeed, CTRW and the relation (28) produce correct dependence of the diffusion coefficient $D(c) \sim c^{4/3} \sim n^{4/3}$ for dipole transport in continuum media approximation, but, according to modern knowledge [31], in the problem of hopping conductivity $D \sim \exp(-\kappa_p / \varepsilon_c^{1/3})$, where $\varepsilon_c = \frac{4\pi}{3} n r_0^3 \ll 1$ and the constant $\kappa_p \sim 1$ is produced by the percolation theory, while CTRW gives other parametric dependence $D = D_{SL} \sim \exp(-\kappa_{SL} / \varepsilon_c^{1/2})$ with $\kappa_{SL} \sim 1$ [2, 17] (see the Appendix).

It should be noted, that all numeric parameters, indicated in this article for dipole transport, are valid for simplified transition rates (7) only. Results for more realistic description can be found in Refs. [4, 28].

Eqs. (1) and (7) are popular in explaining of dipole transport even in the systems, where the conditions of their applicability, indicated after the Eq. (7) are nor fulfilled. It is natural, that, for these systems, results of the CTRW can be applicable for rough qualitative estimations only.

Appendix. Diffusion coefficients in CTRW theory

Short derivation of the diffusion coefficient for the conductivity problem in continuum media approximation is, probably, absent in existing literature. Therefore, we will give it below.

According to relations (14) and (15) (or (26) and (27)) the diffusion coefficient in the Sher-Lax theory is

$$D_{SL} = \frac{1}{2d} \int d^d x x^2 N_{x_0}(\lambda = 0) = \frac{n}{2d} \int d^d x x^2 w_{x_0} \int_0^\infty dt \exp(-w_{x_0} t) Q(t) / \int_0^\infty dt Q(t) \quad (A1)$$

or

$$D_{SL} = \frac{n}{2d} \int_0^\infty dt Q(t) \int d^d x x^2 w_{x_0} \exp(-w_{x_0} t) / \int_0^\infty dt Q(t). \quad (A2)$$

Here d is dimension of the space and the continuum media approximation is applied.

For dipole transport and $d=3$ we have, according to (16), $Q(t) = \exp(-(\beta t)^{1/2})$, and $\int_0^\infty dt Q(t) = 2/\beta$. Then using (A2) and substituting $x = y(w_0 r_0^6 t)^{1/6}$ we obtain

$$D_{SL} = \frac{n}{3\beta} \int \frac{d^3 y}{y^4} \exp(-1/y^6) \int_0^\infty \frac{dt}{t} Q(t) (w_0 r_0^6 t)^{5/6}$$

that directly produces the relation (28), which is exact in continuum media approximation. Similar result was found in Ref. [18].

Calculations for hopping conductivity with $w_{x_0} = w_0 \exp(-x/r_0)$ are much more complex. In order to obtain the main approximation for $w_0 t \gg 1$ we can substitute [2]

$$1 - \exp(-w_{x_0} t) = \mathcal{G}(w_{x_0} t > 1). \quad (A3)$$

As a consequence, in continuum media approximation,

$$Q(w_0 t \gg 1) = \exp\left(-n \int d^3 x (1 - e^{-w_{x_0} t})\right) = \exp\left(-\frac{4\pi}{3} n r_0^3 \ln^3(w_0 t)\right). \quad (A4)$$

For reasonable approximation at all $w_0 t$ this relation can be written as

$$Q(t) = \exp(-\varepsilon_c \ln^3(1 + w_0 t)) \quad (A5)$$

that it is correct in main order for small $\varepsilon_c = \frac{4\pi}{3} n r_0^3 \ll 1$. Therefore

$$\int_0^\infty dt Q(t) = \int_0^\infty dt \exp(-\varepsilon_c \ln^3(1 + w_0 t)) = \int_0^\infty \frac{ds}{w_0} \exp(s - \varepsilon_c s^3) = \int_0^\infty \frac{dy}{\varepsilon_c^{1/2} w_0} \exp\left(\left(y - y^3\right) / \varepsilon_c^{1/2}\right). \quad (A6)$$

where $s = \ln(1 + w_0 t)$. Last integrand has sharp maximum at $y = s \varepsilon_c^{1/2} = \varepsilon_c^{1/2} \ln(1 + w_0 t) = 3^{-1/2}$, that corresponds to exponentially large $w_0 t$, justifies preceding approximations and produces (with exponential accuracy)

$$\int_0^\infty dt Q(t) \sim \frac{1}{w_0} \exp\left(2 / \left(3^{3/2} \varepsilon_c^{1/2}\right)\right). \quad (A7)$$

The preexponent is written here to obtain correct dimensional dependence only.

It is evident that

$$n \int d^3 x w_{x_0} \int_0^\infty dt Q(t) \exp(-w_{x_0} t) = - \int_0^\infty dt \frac{d}{dt} Q(t) = 1. \quad (A8)$$

From other side

$$\begin{aligned} n \int d^3 x w_{x_0} \int_0^\infty dt Q(t) \exp(-w_{x_0} t) &= n \int_0^\infty dt Q(t) \int d^3 x w_{x_0} \exp(-w_{x_0} t) = \\ &= \varepsilon_c w_0 \int_0^\infty dt Q(t) \int dy^3 \exp(-y - w_0 t e^{-y}). \end{aligned} \quad (\text{A9})$$

Last integrand has sharp maximum at $y = y_0(t) = x_0(t) / r_0 = \ln w_0 t \approx \ln(1 + w_0 t)$, and asymptotic expansion in small parameter ε_c near this extremum will restore the identity (A8). Calculation of the diffusion coefficient according to the relation (A2) will produce additional factors x^2 and y^2 in the last integrands of (A9), which can be substituted by $x_0^2(t)$ and $y_0^2(t)$ correspondingly. As a result,

$$\begin{aligned} n \int d^3 x x^2 w_{x_0} \int_0^\infty dt Q(t) \exp(-w_{x_0} t) &= n \int_0^\infty dt Q(t) \int d^3 x x^2 w_{x_0} \exp(-w_{x_0} t) = \\ &= n w_0 \int_0^\infty dt Q(t) \int d^3 x x^2 \exp(-x / r_0 - w_0 t e^{-x/r_0}) \approx \\ &\approx r_0^2 \int_0^\infty dt Q(t) \ln^2(1 + w_0 t) n \int d^3 x w_{x_0} \exp(-w_{x_0} t) = -r_0^2 \int_0^\infty dt \ln^2(1 + w_0 t) \frac{d}{dt} Q(t) = \\ &= -r_0^2 \int_0^\infty \left(d e^{-\varepsilon_c \ln^3(1+w_0 t)} \right) \ln^2(1 + w_0 t) \sim r_0^2 \varepsilon_c^{-2/3}. \end{aligned} \quad (\text{A10})$$

Therefore, according to the relations (A2), (A7) and (A10), we obtain that for hopping conductivity

$$D_{SL} \sim r_0^2 w_0 \exp\left(-2 / \left(3^{3/2} \varepsilon_c^{1/2}\right)\right). \quad (\text{A11})$$

Similar result can be found in Refs. [2, 17].

The relation (A11) is written with exponential accuracy only, that is sufficient to clarify its sharp contradiction with the result of the percolation theory.

References

1. Dzheparov F.S., Lundin A.A. *Sov. Phys. JETP* **48**, 514 (1978)
2. Scher H., Lax M. *Phys. Rev. B* **7**, 4491, 4502 (1973)
3. Gochanour C.R., Andersen H.C., Fayer M.D. *J. Chem. Phys.* **70**, 4254 (1979)
4. Abov Yu.G., Gulko A.D., Dzheparov F.S., Stepanov S.V., Trostin S.S. *Phys. Part. Nucl.* **26**, 692 (1995)
5. Abragam A. *Principles of Nuclear Magnetism*, Oxford University Press, Oxford (1961)
6. Abragam A., Goldman M. *Nuclear Magnetism: Order and Disorder*, Clarendon Press, Oxford (1982)
7. Salikhov K.M., Semenov A.G., Tsvetkov Yu.D. *Elektronnoe spinovoe echo i yego primeneniya*, Nauka, Novosibirsk (1976) (*in Russian*)
8. Atsarkin V.A., Dzheparov F.S. *Z. Phys. Chem.* **231**, 545 (2017)
9. Montroll E.W., Weiss G.H. *J. Math. Phys.* **6**, 167 (1965)
10. Ashurov M.Kh., Basiev T.T., Burshtein A.I., Voron'ko Yu.K., Osiko V.V.. *JETP Lett.* **40**, 841 (1984)
11. Gapontsev V.P., Dzheparov F.S., Platonov N.S., Shestopal V.E. *JETP Lett.* **41**, 561 (1985)
12. Dzheparov F., Gul'ko A., Heitjans P., L'vov D., Schirmer A., Shestopal V., Stepanov S., Trostin S. *Physica B* **297**, 288 (2001)

13. Hughes B.D. *Random Walks and Random Environment*, Clarendon Press, Oxford (1995)
14. Kutner R., Masoliver J. *Eur. Phys. J. B* **90**, 50 (2017)
15. Dzheparov F.S. *Radiospektroskopiya (Perm)* **13**, 135 (1980) (*in Russian*)
16. Dzheparov F.S. *Sov. Phys. JETP* **72**, 546 (1991)
17. Butcher P.N. *J. Phys. C* **7**, 879 (1974)
18. Vugmeister B.E. *Phys. Stat. Sol. (b)* **76**, 161 (1976)
19. Haan S.W., Zwanzig R. *J. Chem. Phys.* **68**, 1879 (1978)
20. Klafter J., Silbey R. *Phys. Rev. Lett.* **44**, 55 (1980)
21. Dzheparov F.S., Smelov V.S., Shestopal V.E. *JETP Lett.* **32**, 47 (1980)
22. Bodunov E.N., Malyshev V.A. *Sov. Phys. Solid State* **26**, 1804 (1984)
23. Dzheparov F.S., Shestopal V.E. *Theor. Math. Phys.* **94**, 345 (1993)
24. Huber D.L., Hamilton D.S., Barnett B. *Phys. Rev. B* **16**, 4642 (1977)
25. Dzheparov F.S., Lvov D.V., Shestopal V.E. *J. Supercond. Novel Magn.* **20**, 175 (2007)
26. Dzheparov F.S. *JETP Lett.* **82**, 521 (2005)
27. Dzheparov F.S., Gulko A.D., Ermakov O.N., Lyubarev A.A., Stepanov S.V., Trostin S.S. *Appl. Magn. Reson.* **35**, 411 (2009)
28. Abov Yu.G., Gulko A.D., Dzheparov F.S., Ermakov O.N., Lvov D.V., Lyubarev A.A. *Phys. At. Nucl.* **77**, 682 (2014)
29. Godzik K., Jortner J. *J. Chem. Phys.* **72**, 4471 (1980)
30. Franchi D.S., Loring R.F., Mukamel S. *J. Chem. Phys.* **86**, 6419 (1987)
31. Zvyagin I.P. *Transport Phenomena in Disordered Semiconductors*, MSU Publ., Moscow (1984) (*in Russian*)